39 N. 5th

An,

Inaugural Dissertation.

Passed March 19
1824
W. E. H
Dean

on,

Hydrocephalus Acutus.

By Cuthbert. T. Green. of Maryland
Licentiate in medicine of Md.
and Member of the Philad.
Medical Society.

Wil

Sir.

in Sei
expos
teid at
dies us
vicke
attention
der Th

To,

William D. Brinckle. A.M. M.D.

Sir.

It affords me much pleasure in seizing on the first opportunity, to express my gratitude for your unremitted attention, while prosecuting my studies under your direction. I am likewise much indebted to you for other attentions equally dear: for which I render this tribute to your friendship.

The Author.

Dissertation 8.

From a consideration of the immense fund of talent displayed in the productions of some of the ancient gentlemen of our profession, I am not a little astonished that hydrocephalus, a disease so pervading in its ravages on the young and rising part of the community, was not noticed as a seperate disease until so late as the year 1768. At which time, it was taken minutely into consideration by Doctor Whytt, who has, I conceive, marked out the course which is to direct us to its correct pathology. Since that period, however, it seems to have at-

tracted

Practice t
midable
Hydroc
ds affect
limited
ers mo
ates, a
children
often yea
non sie
lects; thou
his medi
struggling
respenge
most p
the disea
of myself
delicate
faint, an

tracted that attention, which its formidable nature necessarily demands.

Hydrocephalus, like other dangerous affections, is not by any means limited in its sphere of action. It occurs more particularly in warm climates, and in the mildest seasons. Children from the age of one, to nine or ten years, appear to be the most common victims which this malady selects; though we sometimes see the "vis medicatrix naturæ" in the adult, struggling with this almost certain messenger of death. It is the opinion of most physicians who have noticed the disease, and indeed I have observed it myself, that families, or persons, of a delicate constitutions, scrophulous taint, and more than ordinary brilli-

ancy

may of
Ba
ern the
petable
pecting
formerly
Darwin
supposed
sical Ma
he pres
propriet
natory
ting lis
ate Sen
ly. It is
he mem
phale
his idea
medio

away of intellect, are most subject to it.

Various and contradictory have been the opinions, entertained by respectable practitioners of medicine respecting the nature of this disease. Formerly, physicians as Fothergill, Darwin, Withering, Whytt, Smyth, &c supposed it to be an affection of a dropsical nature; while the physicians of the present day, say it may with more propriety be classed with the inflammatory diseases; and disclaim any thing like an idea of its being a disease similar in any respect to a dropsy. It is to the work of Dr Cheyne, that the new views of the pathology of hydrocephalus may be traced, and it is his ideas that no small share of the medical men of the present day

have a
first g
the a
mente
process
case t
estion,
from t
that t
es fro
the eff
ter aen
ed ac
mina
fact,
that c
this a
ude,
the n

have adopted; without, I am persuaded, first giving them due degree of reflection. The author, whose name I have just mentioned, in his pathology of hydrocephalus says, "That in this disease there is produced a venous congestion, in addition to and arising from the increased arterial action: that the effusion of serous fluid arises from this venous congestion; that the effusion has a tendency to counteract the baleful effects of the increased action, and retard the fatal termination of the disease, is evident in fact, I think it no harsh opinion, that death would occur earlier in this disease, did not the fluid exude, and thus continue to the brain the necessary degree of support"

hence it
proposed
on into
than the
ce, to ag
the existin
this he
source
disposed
absurd.
question
flamm
ble der
view the
a drop
ble con
ty of the
tion. In
by the h

Hence it may be seen that this author is possessed of an opinion, that the effusion into the ventricles is nothing more than the effect, and no way calculated, to aggravate, but rather to mitigate the existing morbid action:-an opinion of this kind although emanating from a source so respectable I am at one disposed to pronounce as inconceivably absurd. That it is produced by a venous congestion succeeded by an increased inflammatory action, no one, I think, will deny. But I am disposed to view the last stages of this disease as of a dropsical nature, and am as well convinced that the uniform fatality of the disease is owing to that action. In this opinion, I am supported by the high authority of Whytt, who

when
immed
dy is th
parts
throw
than
As
of hydr
cor-d a
ject
endea
errors
to esta
optic
1st If
there
ing f
centre
brain

when speaking of this disease, says "the immediate cause of every kind of dropsy is the same; viz such a state of the parts as makes the exhalent arteries throw out a greater quantity of fluid than the absorbents can take up"

As Dr Cheyne's ideas of the nature of hydrocephalus are not in exact accordance with the views on the subject entertained by myself, I shall endeavour to point out some of the errors advanced by him, as well as to establish those which I have adopted as most correct.

1st It is an incontrovertable fact, that there is little or no space for containing fluids in the divisions called ventricles, in a healthy state of the brain; consequently, this effusion into

the mans
extrahe
iting cau
tile mor
not the c
ascribe o
ion take
Cheyne k
his work
le disea
in this M
persons
les ascite
ime, from
the wa
Cheyne's
clus of the
hearts;- fo
consider

"the mansion of life and sensation" as extraneous matter, acts as a re-exciting cause, and renders the disease still more ungovernable. If this be not the case, how can practitioners describe so accurately when this effusion takes place, or why does Dr Cheyne himself, in another part of his work, pronounce it an incurable disease when it does terminate in this manner. The immediate death of persons who have been labouring under ascites for a considerable length of time, from the sudden evacuation of the water, does not strengthen Dr Cheyne's theory in regard to the stimulus of the waters on the adjacent parts;- particularly if we take into consideration the vast difference

of other
of the
3rd Th
lied t
a cons
which
3rd Th
to thos
suppo
costit
mue
of the
respo
colle
of a
the e
citan
pati
has-

of structure, as well as the functions of the parts, implicated in the two diseases.
2nd That it is a dropsy or nearly allied to it, is more than probable from a consideration of the common cause which produces both, viz congestion.
3rd That the symptoms are similar to those that occur in dropsy, as a suppression of urine, of perspiration, costive state of the bowels, hot dry skin, much thirst, paralysis of one side of the head, of one arm, difficult respiration &c symptoms taken collectively are certainly an evidence of a dropsical action. As a proof that the effusion does not act as an irritant or hasten on the death of the patient, it has been stated, that death has taken place after a development.

mes
to hy
con,
rum
cled.
not
esph
kind
ged,
tesist
stag
far o
the
those
mm
A
flam
pect
proof

ment. of the symptoms peculiar to hydrocephalus. and on dissection, nothing like a collection of serum has been found in the ventricles. In such cases, I think it could not with propriety be termed hydrocephalus. as in all terminations of the kind, we have but one of the three stages, which are mentioned as characteristics of the disease, which is the stage of excitement.- in which as far as my limited experience goes the symptoms resembles very much those of an ordinary attack of inflammation of the brain.

As to the theory of its being an inflammation peculiar in any respect, I have seen no satisfactory proof. I ask, is not an effusion into

the be
of a h
mucu
ter in
an in
of the

H
fied
it rus
death
it is
a pum
form
divid
ded
atie
appe
eate
one, a

the ventricles of the brain as the effect of a high degree of inflammation, as much a dropsy, as a collection of water in the thorax as the consequence of an inflammatory action in some one of the viscera of that cavity?

Hydrocephalus is rather diversified in its mode of attack. Sometimes it runs its course and terminates in death in a few days: and sometimes it is more mild in its assault, and assumes a tedious and protracted form; consequently, it calls for some division, and has by some been divided into the Idiopathic and Symptomatic, from the circumstance of its appearing either as an original disease, or as the result of a preceding one, at first quite different. Each of

their fo
ced int
said t
attentio
with th
1st The
ity.
2nd The
tose th
3rd The
Bu
stages
4th The
incipi
are nu
al rem
no di
cutty
discr

these forms have again been divided into three stages, which are said to be known by particular attention to the state of the pulse with other symptoms. They are called
1st The stage of increased sensibility.
2nd The stage of torpor, or comatose stage.
3rd The convulsed stage.

But in very many cases these stages are not well defined.

The symptoms which mark the incipient stage of hydrocephalus are numerous. Indeed it is a general remark by practitioners, that in no disease is there a greater difficulty of laying down a general description of symptoms, than in

this; so p
ons then
myself a
that are
onetal
It begi
bor, wit
domen
face, ho
oma, w
nce in
ermina
the suffe
en, howe
idsable
the atta
ith and
hastice
while the

this; so protiform are the symp
toms themselves. I shall content
myself with delineating those only
that are most prominent, and
general in their appearance.

It begins with a slight degree of
fever, with a pain in the head and
abdomen, sick stomach, flushed
face, hot dry skin, delirium, and
coma, which make their appear-
ance in rapid succession and
terminate the existence of the lit-
tle sufferer in a few days; very of-
ten, however, the patient for a con-
siderable length of time previous to
the attack becomes languid, peu-
vish and uncomfortable, without any
particular complaint. After a
while the appetite is impaired, stom

ach b
iting
sometim
portion
an in
watch
tute th
In
not e
climat
tes upt
throu
child
or put
his he
pulse
mata
the pe
now

ach becomes irritable, with a vomiting of bile, constipated bowels (though sometimes he purges clayey or black fœtid stools) pain in the head with an intolerance of light and sounds, watching and quick pulse constitute the symptoms of the first stage.

In the second, the patient is not easily aroused, has a great inclination to sleep, but is much interupted with sharp pains shooting through his head; when in bed, the child is constantly rolling about, or putting his hands up towards his head, his pupil is dilated, pulse slow, suffusion of the adnata, and the eye which was in the first stage morbidly sensible, now loses its lustre and expression-

The th
reasedo
hurrie
ulsion
versale
practi
death
Noth
than th
of the p
ache a
of unti
which
the ex
action
estime
tons
child
The e

The third stage is known by the increased arterial action, strabismus, hurried breathing, stupor and convulsions; symptoms which are universally regarded by the discerning practitioner as the harbingers of death.

Nothing however can be more irregular than the progress and development of the foregoing symptoms; the head ache is very often not complained of until effusion has taken place, which is a very striking proof of the existence of the inflammatory action without much pain. At other times, the most alarming symptoms will suddenly remit, and the child at the same time in eminent danger. The eye has heretofore been looked

upon a
criterio
some f
founda
I s
vations
hydro c
most p
is like
of the
is in c
of bloo
the bra
the bo
ble to
the me
drink
irritat
rect. A

upon as one of the most unerring criterions, but like the pulse, in some particular cases, it will be found very fallacious.

I shall now make a few observations on the symptomatic form of hydrocephalus. This is by far the most frequent in its appearance and is likewise the most manageable of the two kinds. It appears that there is in childhood a greater supply of blood to the digestive organs, and the brain, than to any other parts of the body. These two organs are liable to numerous derangements, from the morbid stimulation of food and drink on the one part, and mental irritation on the other. From the direct sympathy existing between the

two or g
disease
into a
then, a
the br
with a
alimen
tion or
tions o
cera, b
that of
all ca
duce
liver; s
the con
this m
is pro
ter m
tants

two organs. one cannot continue long diseased without drawing the other into a similar state. This disease then, almost always occurs from the brain morbidly sympathising with a disordered state of the alimentary canal; or from a congestion or an interruption of ~~the~~ functions of some of the abdominal viscera, but more particularly with that of the liver; Currie supposed all cases of hydrocephalus were produced by previous irritation of the liver; such ideas I think rather pass the confines of reason. In what way this morbid sympathetic influence is propagated, is not for me to determine in this place. That irritants in the primæ viæ bring on this

disease
proof th
most c
licence
medic
Symp
on in a
to an
appear
deran
the co
domin
dent.
be pre
of the
The
use the
gnor,
dulln

disease, we need go no further for proof than the fact, of some of the most alarming symptoms being relieved by a recourse to purgative medicines.

Symptomatic hydrocephalus comes on in a slow insidious manner, and to an inattentive observer it would appear that there was little or no derangement taking place:- but in the course of a few days, some abdominal derangement will be evident. The disease at this time may be prevented by a prompt removal of the primary source of irritation.

The symptoms which characterize this form of the disease, are languor, an unhealthy countenance dullness of the eye, skin hot and

dry, bowel
and tong
the stools
white cla
now be c
sore: the
and the
to the gen
children
cision of
prehensi

brain
hydroc
gid
membr
lining
sidera
ening o

dry, bowels torpid appetite irregular and tongue white. If the bowels be open, the stools here as in the other form, are white clayey or bilious. The abdomen now becomes somewhat swollen and sore: the pulse a little accelerated and the head is likewise drawn into the general disorder. Dr Yeates says "children at this period evince a precision of ideas and quickness of apprehension much beyond their years"

The appearances of the brain of children who die of hidrocephalus are an engorged state of the vessels of the membrane spread over, and lining the ventricles, and considerable adhesions as well as thickening of this structure will likewise

be observ-
mon of al
the effusi
ventricle
brain, or
mater. T
be the effe
matory a
the part,
of nature
overloade
tem. The
ately prov
ces both
loci; Ma
ny others
injury b
proper
I m

be observed. But the most common of all morbid appearances is the effusion of water either in the ventricles, or between the lobes of the brain, or between the dura and pia mater. This is generally believed to be the effect of the increased inflammatory action of the blood vessels of the part, and appears to be an effort of nature, for the present relief of the overloaded state of the vascular system. This very effusion though ultimately proves fatal in its consequences both by compression and "error loci", Nature in this case, as in many others, produces irremediable injury by relieving herself at an improper place.

I now come to treat of the mode

of cure,
tient lab
lus. we
in our
ries, not
toms the
ous sta
that we
the kind
very diff
toms wh
always
by the o
is suffi
The
the first
rary ac
which
such as

of cure, when first called to a patient labouring under hydrocephalus, we should be very particular in our examinations and enquiries, not only in regard to symptoms then existing, but to the previous state of the child, in order that we may if possible designate the kind, an object in some cases very difficult to effect, as the symptoms which I have stated are not always well developed consequently the diagnosis in many cases is difficult.

The treatment to be adopted in the first stage of this disease should vary according to the symptoms which present themselves. If they are such as were mentioned as characteristic

wishing a
opathic fo
the natur
a mom
ent, whic
discrimin
ded some
next pes
these cal
attention
constipat
liver, its
secretions
the proper
omment
disease, L
propriety
judicio
of our su

terising an incipient attack of the idiopathic form; no one conversant with the nature of the disease would hesitate a moment to recur to the use of the lancet, which should be used freely but with discrimination. After having afforded some relief by this practice, our next resource is from purgatives. Of these calomel I think claims our first attention. Its efficacy in overcoming constipation, its peculiar power on the liver, its promptness in altering the secretions of the stomach, bowels &c are the properties which particularly recommend its employment in this disease. Too much cannot be said of the propriety of purging here, as it is on the judicious exhibition of them, that much of our success depends. Calomel

combined with
administered
cure copious
ed, will be fo

But if
gent, the jug
tay should be
considerable
with this m
should be sha
or perhaps le
purpose, un
as they detra
and are not
ing symptom
cupping is
course when
will sometimes
effects. Non s

combined with gamboge or jalap, and administered in such doses as to procure copious evacuations, daily repeated, will be found very beneficial.

But if cerebral irritation be urgent, the jugular vein or temporal artery should be opened and blood let to a considerable extent. In conjunction with this mode of treatment, the head should be shaved and cups applied; or perhaps leeches would serve a better purpose, under this circumstances, as they detract blood with more ease and are not attended with the distressing symptoms which the operation of cupping is found to produce. This course when vigorously persisted in will sometimes be productive of very salutary effects. Nor should the practitioner let

the adv-an
plicted as
part on wh
its force, a
of attention
stances con
to a belief t
disturban
companied
trial actio
to disconte
and resort
as pour-de
out of ethe
and water

As yet
the use of
plication
important

the advantage thus gained be neglected as the bowels, as well as the part on which the disease spends its force, require the utmost degree of attention. If however some circumstances continue which would lead to a belief in the continuance of the disturbance of the encephalon, accompanied with a diminution of arterial action, it would be advisable to discontinue the use of the lancet, and resort to cold applications such as powdered ice or cloths wrung out of ether and water or vinegar and water.

As yet I have said nothing of the use of blisters. the well timed application of which constitute very important remedies in the manag

ment of my
be applied t
the forehead
very often
discharge,
may effect
be dissed u
n if there a
ast to hea
applieds.

A good
the powers
But as to i
other disea
tain stre
bled to s
ers very h
taken pla
n of arresti

ment of hydrocephalus. they should be applied to the nape of the neck, on the forehead, or all over the heads. It is very often necessary to keep up the discharge, to which alone the salutary effect is due, The blister should be dressed with mercurial ointment or if there appears a disposition in the part to heal the blister might be re-applieds.

A good deal has been said of the powers of digitalis in this disease But as to its influence in this or any other disease requiring depletion I entertain strong doubts, Nor am I disposed to value its remediate powers very highly after effusion has taken place. that it possesses the power of arresting the circulation no one

vide position a
y contra tha
iciently imp
t application
ppe it is no
at I oppose
rience in
is or any oth
mboldened
t fall by m
ken into w
ulli; I mak
the above ac
d drawn fr
t use
When eff
hole reliar
hudden int
the system

will pretend to deny, but I strenuously contend that this property is not sufficiently important as to recommend its application in this disease. I must confess it is not with a little diffidence that I oppose the opinions of men of experience in regard to the operation of this or any other medicine, but being emboldened by the liberal principles let fall by more than one of the gentlemen into whose hands this essay is to fall; I make no hesitation in giving the above as my candid belief, which is drawn from some experience in its use.

When effusion takes place our whole reliance is to be placed on the sudden introduction of mercury into the system. This action should be in-

duced

duced by the
calomel in
mercurial
y considera
on such cir
than it is
If it shoul
d symptoms.
sympathet
the persuad
dear the wh
fensive co
hich, calom
much arter
must be w
f such an
The emacia
nutritious t
continuan

duced by the prompt exhibition of calomel in large doses, together with mercurial frictions, employed to a very considerable extent, the ointment under such circumstances should be stronger than it is ordinarily made.

If it should come on with the train of symptoms mentioned as indicating a sympathetic affection, the course then to be pursued is self evident: which is to clear the whole line of intestines of its offensive contents, with a preparation, of which, calomel is the base. If there be much arterial excitement the lancet must be unsheathed, but not carried to such an extent as in the idiopathic. The evacuations themselves are the best criterions to direct us in regard to the continuance of the purgative course;

After eff
ith in prom
n one, an
owledge o
ntents in
ostly take
nce of the
ware that
had arisen.
gainst it, it
hat the Sp
ne very eg
n imitated
ebral resp
then let us
which is n
mopes in
make the c
known to be

After effusion has taken place all unite in pronouncing the case a desperate one, an opinion drawn from a knowledge of the inactivity of the absorbents in the part where the effusion mostly takes place. From the prevalence of the foregoing opinion I am aware that much mischief at times has arisen. In order then to guard against it, it should be borne in mind that the symptoms of actual effusion are very equivocal, and may be closely imitated by a turgescence of the cerebral vessels; From this circumstance then let us learn a practical precept. which is not to discontinue the remedies proper in effusion until symptoms make their appearance which we know to be the precursors of death;

Of howe
sick under
some symptom
which precede
termination
part is nec
where a gre
than in this
in a very de
ence of so
subsiding
supporting
effect this
of tonic
ing diet
exercise and

If however the child does not sink under the violence of the disease, and symptoms manifest themselves which predict a more favourable termination, much care on our part is necessary; as in no disease is there a greater liability to a relapse than in this. The child is mostly left in a very debilitated state on the subsidence of so formidable a disease. Our business then is to use every means of supporting the strength of the child. To effect this we must resort to the use of tonics and a mild and nourishing diet in conjunction with gentle exercise and change of air.

Cant effusion reach b. 2

~~Ste~~ Not dropsy from b. asterin.